

THE USE OF THE ENZYME-LINKED IMMUNOSORBENT ASSAY AND  
IMMUNOSORBENT ELECTRON MICROSCOPY FOR THE DETECTION AND  
CHARACTERIZATION OF CASSAVA COMMON MOSAIC VIRUS

*(Usage du Test ELISA et de l'immunoélectromicroscopie pour  
détecter et caractériser une souche de la mosaïque commune du manioc)*

B. PINEDA, A.C. VELASCO and B.L. NOLT

CIAT Cassava Program  
Apdo. Aéro 6713  
CALI, COLOMBIA

SUMMARY

The double-antibody sandwich form of the enzyme-linked immunosorbent assay (ELISA) provided a rapid and sensitive means for the detection of cassava common mosaic virus (CCMV) in cassava (*Manihot esculenta* Crantz). The ELISA procedure is particularly useful in a large-scale CCMV indexing program because many more plants or plantlets can be assayed per day than by graft indexing or bioassays. The horseradish peroxidase : O = phenylenediamine enzyme-substrate combination is a sensitive signal for detecting CCMV. The immunosorbent electron microscopy (ISEM) technique was used to serologically compare a Colombian isolate with CCMV and other members of the potexvirus group by observing the number of virus particles selectively absorbed by different potexvirus antisera. The ISEM method also increased the number of virus particles adsorbed on the surface of the electron microscope grid available for observation and measurement.

RESUME

*Le test immunoenzymatique à double anticorps "ELISA" fournit une méthode rapide et sensible pour détecter la mosaïque commune du manioc (CCMV) dans les plants de Manioc.*

*Le test ELISA est particulièrement utile pour des tests d'indexation à grande échelle, car il permet d'examiner beaucoup plus de plantes par jour que par les tests utilisant le greffage ou les hôtes différentiels. La combinaison enzyme-substrat peroxydase du radis noir - orthophénylène-diamine peut servir de signal particulièrement sensible pour détecter CCMV. L'immunoélectromicroscopie (ISEM) a été utilisée pour comparer sérologiquement une souche colombienne de CCMV et*

*d'autres membres du groupe "potexvirus" en observant le nombre de particules virales sélectivement adsorbées par les antisérums correspondant aux divers potexvirus. Cette méthode permet aussi d'augmenter le nombre de particules virales fixées sur les grilles pour observation et mensuration au microscope électronique.*

The double antibody sandwich form of the enzyme-linked immunosorbent assay (ELISA) provided a rapid and sensitive means for the detection of cassava common mosaic virus (CCMV) in cassava (*Manihot esculenta* Crantz). The ELISA procedure is particularly useful in a large-scale CCMV indexing program because many more plants or plantlets can be assayed per day than by graft indexing or bioassays. The horseradish peroxidase : O = phenylenediamine enzyme-substrate combination is a sensitive signal for detecting CCMV. The immunosorbent electron microscopy (ISEM) technique was used to serologically compare a Colombian isolate with CCMV and other members of the potexvirus group by observing the number of virus particles selectively absorbed by different potexvirus antisera. The ISEM method also increased the number of virus particles adsorbed on the surface of the electron microscope grid available for observation and measurement.

Cassava common mosaic virus (CCMV), a plant potexvirus with elongated rod-shaped particles, is present in cassava from many Latin American countries. The field incidence of CCMV is usually low however the yield of infected plants can be reduced by as much as 60 per cent. Although there are no reported vectors of CCMV this virus is readily mechanically transmitted and efficiently disseminated in infected planting material. The risk of introducing CCMV into new areas via infected vegetative material is high therefore sensitive virus detection methods are required for indexing cassava germplasm for the presence of CCMV. The double antibody sandwich form of the enzyme-linked immunosorbent assay (ELISA) and immunosorbent electron microscopy (ISEM) are available for detecting CCMV in cassava.

The ELISA test is conducted in small wells in a polystyrene or polyvinyl chloride microtiter plate containing 96 sample wells. Sample preparation consists of grinding the plant tissue sample in 20 to 100 volumes of extraction buffer with a mortar and pestle. We found that buffer dilutions greater than 1/10 worked better for cassava tissue. Depending on the available labor up to several hundred plants can be indexed/day with results available within 36 hours. If required many more samples can be processed using commercially available tissue extractors and automated ELISA systems. We routinely use CCMV gamma globulins conjugated to horseradish peroxidase as the detecting antibodies but alkaline phosphatase can also be used. Using ELISA CCMV can be detected in small leaf disks from infected cassava plants or in small leaves from plantlets.

ISEM combines the specificity of serology with electron microscopy. Virus particles are selectively adsorbed or "trapped" on the surface of EM grids pre-coated with related antibody molecules increasing the number available for observation. The number of virus particles is related to the degree of serological specificity between the coating antibody and the virus. The higher the number of particles trapped the closer the serological relationship. This technique was used to determine the serological relationship of a Colombian isolate of CCMV with other members of the plant potexvirus group. The ISEM technique can also provide a rapid virus identification. The presence of CCMV can be determined within several hours.

